

Rebecca



Before JUVÉDERM® treatment



After 2.5 syringes
JUVÉDERM® Ultra (1.2 mL)
JUVÉDERM® Ultra Plus (0.8 mL)



Before JUVÉDERM® treatment



After 2.5 syringes
JUVÉDERM® Ultra (1.2 mL)
JUVÉDERM® Ultra Plus (0.8 mL)



Before
JUVÉDERM® treatment



After 1 syringe
JUVÉDERM® Ultra Plus
(0.8 mL)
(2 weeks after injection)



After 2 syringes
JUVÉDERM® Ultra (0.8 mL)
JUVÉDERM® Ultra Plus (0.8 mL)
(2 weeks after injection of second syringe)



After 2.5 syringes
JUVÉDERM® Ultra (1.2 mL)
JUVÉDERM® Ultra Plus (0.8 mL)
(2 weeks after injection of third syringe)

Unretouched photographs. Individual results may vary.

In the United States, JUVÉDERM® injectable gel is indicated for correction of moderate to severe facial wrinkles and folds (such as nasolabial folds).

JUVÉDERM® Important Safety Information

JUVÉDERM® injectable gel is indicated for correction of moderate to severe facial wrinkles and folds (such as nasolabial folds). Side effects were usually mild to moderate, lasting 7 days or less, and included temporary injection-site reactions such as redness, pain, firmness, swelling, and bumps. JUVÉDERM® is not for people with severe allergies. For more information, please click on the About Safety link at www.juvederm.com or call the Allergan Product Support line at 1-877-345-5372. JUVÉDERM® injectable gel is available by prescription only.